arXiv:1302.4222v1 [math.CA] 18 Feb 2013

THE RADIUS OF CONVEXITY OF NORMALIZED BESSEL FUNCTIONS OF THE FIRST KIND

ÁRPÁD BARICZ AND RÓBERT SZÁSZ

Abstract. In this paper we determine the radius of convexity for two kind of normalized Bessel functions of the first kind. In the mentioned cases the normalized Bessel functions are starlike-univalent and convex-univalent, respectively, on the determined disks.

1. Introduction

Let $\mathbf{D}(z_0,r)=\{z\in\mathbb{C}:|z-z_0|<r\}$ denote the open disk centered in z_0 and of radius $r>0$, and let $\mathcal{A}$ be the class of analytic functions defined in the open unit disk $\mathbf{D}=\{z\in\mathbb{C}:|z|<1\}$ and having the property that $f(0)=f'(0)-1=0$. Recall that a function $f\in\mathcal{A}$ belongs to the class $\mathcal{K}$ of convex functions if and only if the image domain $f(\mathbf{D})$ is a convex domain in $\mathbb{C}$, that is, the domain $f(\mathbf{D})\subset\mathbb{C}$ contains the entire line segment joining any pair of its points. It is well-known that the class of convex functions can be characterized as

$$\mathcal{K}=\left\{f\in\mathcal{A}\,\middle|\,\mathrm{Re}\left(1+\frac{zf''(z)}{f'(z)}\right)>0,\ z\in\mathbf{D}\right\}.$$

Moreover, for $\alpha\in[0,1)$ we consider also the class of convex functions of order α defined by

$$\mathcal{K}(\alpha)=\left\{f\in\mathcal{A}\,\middle|\,\mathrm{Re}\left(1+\frac{zf''(z)}{f'(z)}\right)>\alpha,\ z\in\mathbf{D}\right\}.$$

Now let us consider the radius of convexity, and the radius of convexity of order α of the function f

$$r^c(f)=\sup\left\{r\in(0,\infty)\,\middle|\,\mathrm{Re}\left(1+\frac{zf''(z)}{f'(z)}\right)>0,\quad z\in\mathbf{D}(0,r)\right\}$$

and

$$r^c_\alpha(f)=\sup\left\{r\in(0,\infty)\,\middle|\,\mathrm{Re}\left(1+\frac{zf''(z)}{f'(z)}\right)>\alpha,\quad z\in\mathbf{D}(0,r)\right\}.$$

We note that $r^c(f)$ is in fact the largest radius for which the image domain $f(\mathbf{D}(0,r^c(f)))$ is a convex domain in $\mathbb{C}$.

The Bessel function of the first kind of order ν is defined by

$$J_\nu(z)=\sum_{n\geq0}\frac{(-1)^n}{n!\Gamma(n+\nu+1)}\left(\frac{z}{2}\right)^{2n+\nu}.$$

2010 *Mathematics Subject Classification.* 33C10, 30C45.

Key words and phrases. Bessel functions of the first kind; starlike and convex functions; radius of starlikeness; radius of convexity.

In this paper we focus on the following normalized forms

$$f_\nu(z) = (2^\nu \Gamma(\nu+1) J_\nu(z))^{\frac{1}{\nu}} = z - \frac{1}{4\nu(\nu+1)} z^3 + \dots, \ \nu \neq 0,$$

$$g_\nu(z) = 2^\nu \Gamma(\nu+1) z^{1-\nu} J_\nu(z) = z - \frac{1}{4(\nu+1)} z^3 + \frac{1}{32(\nu+1)(\nu+2)} z^5 - \dots,$$

where $\nu > -1$. Observe that $f_\nu, g_\nu \in \mathcal{A}$. We also mention that

$$f_\nu(z) = \exp\left(\frac{1}{\nu} \operatorname{Log}\left(2^\nu \Gamma(\nu+1) J_\nu(z)\right)\right),$$

where Log represents the principal branch of the logarithm, and in this paper every multi-valued function is taken with the principal branch. We also mention that the univalence, starlikeness and convexity of other functions involving the Bessel function of the first kind were studied extensively in several papers. We refer to [**Ba1**, **Ba2**, **Ba3**, **BKS**, **BP**, **Br**, **KT**, **Sz**, **SK**] and to the references therein.

In this paper we make a further contribution to the subject by showing the following new results. The next section contains some preliminary results, and the proofs of Theorems 1 and 2 can be found in section 3.

Theorem 1. *If $\nu > 0$ and $\alpha \in [0,1)$, then the radius of convexity of order α of the function f_ν is the smallest positive root of the equation*

$$1 + \frac{rJ_\nu''(r)}{J_\nu'(r)} + \left(\frac{1}{\nu} - 1\right) \frac{rJ_\nu'(r)}{J_\nu(r)} = \alpha. \tag{1.1}$$

Moreover $r_\alpha^c(f_\nu) < j_{\nu,1}' < j_{\nu,1}$, where $j_{\nu,1}$ and $j_{\nu,1}'$ denote the first positive zeros of J_ν and J_ν', respectively.

Theorem 2. *If $\nu > -1$ and $\alpha \in [0,1)$, then the radius of convexity of order α of the function g_ν is the smallest positive root of the equation*

$$1 + r\frac{rJ_{\nu+2}(r) - 3J_{\nu+1}(r)}{J_\nu(r) - rJ_{\nu+1}(r)} = \alpha. \tag{1.2}$$

Moreover, $r_\alpha^c(g_\nu) < \alpha_{\nu,1} < j_{\nu,1}$, where $\alpha_{\nu,1}$ is the first positive zero of the Dini function $z \mapsto (1-\nu)J_\nu(z) + zJ_\nu'(z)$.

The real number

$$r^*(f) = \sup\left\{ r \in (0,\infty) \middle| \operatorname{Re}\left(\frac{zf'(z)}{f(z)}\right) > 0, \quad z \in \mathbf{D}(0,r) \right\}$$

is called the radius of starlikeness of the function f and it is the largest radius such that $f\left(\mathbf{D}(0, r^*(f))\right)$ is a starlike domain with respect to 0. It is important to mention here that recently, the authors and Kupán [**BKS**] investigated the radius of starlikeness of the functions f_ν and g_ν, and showed that for $\nu > 0$ the radius of starlikeness of f_ν is $j_{\nu,1}'$, while for $\nu > -1$ the radius of starlikeness of g_ν is $\alpha_{\nu,1}$. The above results on radii of starlikeness of f_ν and g_ν for $\nu > 0$ were proved earlier by Brown [**Br**], by using a different approach. For more details we refer to [**BKS**].

2. Lemmas

Lemma 1. *If $a > b > 0$, $z \in \mathbb{C}$ and $\lambda \in [0, 1]$, then for all $|z| < b$ we have*

$$\lambda \operatorname{Re}\left(\frac{z}{a-z}\right) - \operatorname{Re}\left(\frac{z}{b-z}\right) \geq \lambda \frac{|z|}{a-|z|} - \frac{|z|}{b-|z|}. \tag{2.1}$$

Proof. Let us consider the function $u : [b, \infty) \to \mathbb{R}$, defined by

$$u(t) = \operatorname{Re}\left(\frac{z}{t-z}\right) - \frac{|z|}{t-|z|}.$$

Simple computations lead to

$$u(t) = \frac{tx - m^2}{t^2 - 2xt + m^2} - \frac{m}{t-m},$$

$$u'(t) = \frac{2tm^2 - t^2x - xm^2}{(t^2 - 2xt + m^2)^2} + \frac{m}{(t-m)^2},$$

where $z = x + \mathrm{i}y$ and $|z| = m$. Since for $t > 0$ we have

$$u'(t) \geq \frac{2tm^2 - t^2x - xm^2}{(t^2 - 2xt + m^2)^2} + \frac{m}{t^2 - 2xt + m^2} = \frac{(m-x)(t+m)^2}{(t^2 - 2xt + m^2)^2} > 0,$$

it follows that u is an increasing function, and consequently we get $u(a) \geq u(b)$, which is equivalent to

$$\operatorname{Re}\left(\frac{z}{a-z}\right) - \frac{|z|}{a-|z|} \geq \operatorname{Re}\left(\frac{z}{b-z}\right) - \frac{|z|}{b-|z|}. \tag{2.2}$$

On the other hand, it is known (see [**Sz**]) that if $z \in \mathbb{C}$ and $\mu \in \mathbb{R}$ such that $\mu > |z|$, then

$$\frac{|z|}{\mu - |z|} \geq \operatorname{Re}\left(\frac{z}{\mu - z}\right). \tag{2.3}$$

This in turn implies that

$$\lambda \left[\operatorname{Re}\left(\frac{z}{a-z}\right) - \frac{|z|}{a-|z|}\right] \geq \operatorname{Re}\left(\frac{z}{a-z}\right) - \frac{|z|}{a-|z|}.$$

and combining this with (2.2) we get (2.1). □

Lemma 2. [**Wa**, p. 482] *If $\nu > -1$ and $a, b \in \mathbb{R}$, then $z \mapsto aJ_\nu(z) + bzJ_\nu'(z)$ has all its zeros real, except the case when $a/b + \nu < 0$. In this case it has two purely imaginary zeros beside the real roots. Moreover, if $\nu > -1$ and $a, b \in \mathbb{R}$ such that $a^2 + b^2 \neq 0$, then no function of the type $z \mapsto aJ_\nu(z) + bzJ_\nu'(z)$ can have a repeated zero other than $z = 0$.*

Lemma 3. [**Wa**, p. 198] *If $\nu > -1$, then for the Bessel functions of the third kind $H_\nu^{(1)}$ and $H_\nu^{(2)}$ the following asymptotic expansions are valid*

$$H_\nu^{(1)}(w) = \left(\frac{2}{\pi w}\right)^{\frac{1}{2}} e^{\mathrm{i}(w - \frac{1}{2}\nu\pi - \frac{1}{4}\pi)}(1 + \eta_{1,\nu}(w)),$$

$$H_\nu^{(2)}(w) = \left(\frac{2}{\pi w}\right)^{\frac{1}{2}} e^{-\mathrm{i}(w - \frac{1}{2}\nu\pi - \frac{1}{4}\pi)}(1 + \eta_{2,\nu}(w)),$$

where $\eta_{1,\nu}(w)$ and $\eta_{2,\nu}(w)$ are $\mathcal{O}(1/w)$ when $|w|$ is large.

Lemma 4. *Let $\alpha_{\nu,n}$ be the n-th positive root of the equation $J_\nu(z) - zJ_{\nu+1}(z) = 0$. If $\nu > -1$, then the following development holds*

$$\frac{g_\nu''(z)}{g_\nu'(z)} = \frac{zJ_{\nu+2}(z) - 3J_{\nu+1}(z)}{J_\nu(z) - zJ_{\nu+1}(z)} = -\sum_{n\geq 1} \frac{2z}{\alpha_{\nu,n}^2 - z^2}. \tag{2.4}$$

Proof. Let us consider the integral

$$\frac{1}{2\pi \mathrm{i}} \int_{\mathbf{U}} \frac{z}{w(w-z)} \frac{wJ_{\nu+2}(w) - 3J_{\nu+1}(w)}{J_\nu(w) - wJ_{\nu+1}(w)} dw$$

where $\mathbf{U}$ is the rectangle, whose vertices are $\pm a \pm b\mathrm{i}$, $a > 0$, $b > 0$ and z is a point inside the rectangle $\mathbf{U}$ other than a zero of $z \mapsto J_\nu(z) - zJ_{\nu+1}(z)$. Suppose that inside of $\mathbf{U}$ there are m positive and m negative roots of $z \mapsto J_\nu(z) - zJ_{\nu+1}(z)$. Since $(1-\nu)J_\nu(z) + zJ_\nu'(z) = J_\nu(z) - zJ_{\nu+1}(z)$, according to Lemma 2 the zeros of $z \mapsto J_\nu(z) - zJ_{\nu+1}(z)$ are simple and real. The point $w = 0$ is an eliminable singularity. The only poles of the above integrand inside the rectangle are z, $\pm\alpha_{\nu,1}$, $\pm\alpha_{\nu,2}$, $\dots$, $\pm\alpha_{\nu,m}$. The residue at z is

$$\varphi_\nu(z) = \frac{zJ_{\nu+2}(z) - 3J_{\nu+1}(z)}{J_\nu(z) - zJ_{\nu+1}(z)},$$

while the residues at $\pm\alpha_{\nu,n}$ are

$$\frac{z}{\alpha_{\nu,n}(\alpha_{\nu,n} \mp z)}.$$

Here we used the recurrence relation

$$zJ_\nu'(z) = -zJ_{\nu+1}(z) + \nu J_\nu(z)$$

for ν and $\nu + 1$. The residue theorem implies that

$$\begin{aligned} &\frac{1}{2\pi \mathrm{i}} \int_D \frac{z}{w(w-z)} \frac{wJ_{\nu+2}(w) - 3J_{\nu+1}(w)}{J_\nu(w) - wJ_{\nu+1}(w)} dw \\ &\quad = \frac{zJ_{\nu+2}(z) - 3J_{\nu+1}(z)}{J_\nu(z) - zJ_{\nu+1}(z)} + \sum_{n=1}^m \frac{z}{\alpha_{\nu,n}(\alpha_{\nu,n} - z)} + \sum_{n=1}^m \frac{z}{\alpha_{\nu,n}(\alpha_{\nu,n} + z)} \\ &\quad = \frac{zJ_{\nu+2}(z) - 3J_{\nu+1}(z)}{J_\nu(z) - zJ_{\nu+1}(z)} + \sum_{n=1}^m \frac{2z}{\alpha_{\nu,n}^2 - z^2}. \end{aligned}$$

In what follows we show that a and b can be replaced by suitable sequences which increase without limit and in the same time the expression $\varphi_\nu(w)$ remains bounded if $w \in \mathbf{U}$. Since the function $w \mapsto \varphi_\nu(w)$ is an odd function of w, it is sufficient to consider the case $\operatorname{Re} w > 0$. Now, we introduce the notations

$$H_{\nu+k}^{(j)}(w) = \left(\frac{2}{\pi w}\right)^{\frac{1}{2}} e^{(-1)^{j-1}\mathrm{i}(w - \frac{1}{2}(\nu+k)\pi - \frac{1}{4}\pi)}(1 + \eta_{k,j,\nu}(w)),$$

where, as in Lemma 3, for $k \in \{0,1,2\}$ and $j \in \{1,2\}$ the expression $\eta_{k,j,\nu}(w)$ is $\mathcal{O}(1/w)$ when $|w|$ is large. The relation $2J_\nu(w) = H_\nu^{(1)}(w) + H_\nu^{(2)}(w)$ and Lemma 3 lead to $\varphi_\nu(w) = p_\nu(w)/q_\nu(w)$, where

$$\begin{aligned} p_\nu(w) = &(-\mathrm{i})e^{2\mathrm{i}(w - \frac{1}{2}(\nu+1)\pi - \frac{1}{4}\pi)}(1 + \eta_{2,1,\nu}(w)) + \mathrm{i}(1 + \eta_{2,2,\nu}(w)) \\ &- \frac{3}{w}\left[e^{2\mathrm{i}(w - \frac{1}{2}(\nu+1)\pi - \frac{1}{4}\pi)}(1 + \eta_{1,1,\nu}(w)) + (1 + \eta_{1,2,\nu}(w))\right] \end{aligned}$$

and

$$q_\nu(w) = \frac{1}{w}\left[\mathrm{i}e^{2\mathrm{i}(w-\frac{1}{2}(\nu+1)\pi-\frac{1}{4}\pi)}(1+\eta_{0,1,\nu}(w)) - \mathrm{i}(1+\eta_{0,2,\nu}(w))\right]$$
$$-e^{2\mathrm{i}(w-\frac{1}{2}(\nu+1)\pi-\frac{1}{4}\pi)}(1+\eta_{1,1,\nu}(w)) + (1+\eta_{1,2,\nu}(w)).$$

Since $\varphi_\nu(x+\mathrm{i}b)$ tends to i as $b\to\infty$ and the convergence is uniform with respect to $x\in\mathbb{R}$, it follows that $\varphi_\nu(w)$ is bounded on $\{x+\mathrm{i}b | x\in[0,a]\}$ if b is large enough. An analogous argument shows that $\varphi_\nu(w)$ is bounded on the segment $\{x-\mathrm{i}b | x\in[0,a]\}$. It remains to prove that $\varphi_\nu(w)$ is bounded in the case when $w\in\{a+\mathrm{i}y | y\in[-b,b]\}$. In this case we put $a=a_m=2m\pi+\frac{1}{2}(\nu+1)+\frac{3}{4}\pi$ and we get

$$\lim_{m\to\infty}\frac{p_\nu(2m\pi+\frac{1}{2}(\nu+1)+\frac{3}{4}\pi+\mathrm{i}y)}{q_\nu(2m\pi+\frac{1}{2}(\nu+1)+\frac{3}{4}\pi+\mathrm{i}y)} = \frac{\mathrm{i}(1+e^{-2y})}{1+e^{-2y}} = \mathrm{i},$$

and thus it follows that $\varphi_\nu(w)$ is bounded on the segment $\{a+\mathrm{i}y | y\in[-b,b]\}$, if a_m is large enough. Consequently $\varphi_\nu(w)$ is bounded on the perimeter of the rectangle $\mathbf{U}$ if b and a_m tend to infinity. Thus it follows that

$$\lim_{\substack{b\to\infty\\ m\to\infty}}\frac{1}{2\pi\mathrm{i}}\int_{\mathbf{U}}\frac{z}{w(w-z)}\frac{wJ_{\nu+2}(w)-3J_{\nu+1}(w)}{J_\nu(w)-wJ_{\nu+1}(w)}dw = 0,$$

which implies that (2.4) is indeed valid. □

3. Proofs of the Main Results

Proof of Theorem 1. Observe that

$$1+\frac{zf_\nu''(z)}{f_\nu'(z)} = 1+\frac{zJ_\nu''(z)}{J_\nu'(z)}+\left(\frac{1}{\nu}-1\right)\frac{zJ_\nu'(z)}{J_\nu(z)}.$$

Now, recall the following infinite product representations [**OLBC**, p. 235]

$$J_\nu(z) = \frac{\left(\frac{1}{2}z\right)^\nu}{\Gamma(\nu+1)}\prod_{n\geq1}\left(1-\frac{z^2}{j_{\nu,n}^2}\right),\quad J_\nu'(z) = \frac{\left(\frac{1}{2}z\right)^{\nu-1}}{2\Gamma(\nu)}\prod_{n\geq1}\left(1-\frac{z^2}{j_{\nu,n}'^2}\right),$$

where $j_{\nu,n}$ and $j_{\nu,n}'$ are the nth positive roots of J_ν and J_ν', respectively. Logarithmic differentiation yields

$$\frac{zJ_\nu'(z)}{J_\nu(z)} = \nu-\sum_{n\geq1}\frac{2z^2}{j_{\nu,n}^2-z^2},\quad 1+\frac{zJ_\nu''(z)}{J_\nu'(z)} = \nu-\sum_{n\geq1}\frac{2z^2}{j_{\nu,n}'^2-z^2},$$

which implies that

$$1+\frac{zf_\nu''(z)}{f_\nu'(z)} = 1-\left(\frac{1}{\nu}-1\right)\sum_{n\geq1}\frac{2z^2}{j_{\nu,n}^2-z^2}-\sum_{n\geq1}\frac{2z^2}{j_{\nu,n}'^2-z^2}.$$

Now, suppose that $\nu\in(0,1]$. By using the inequality (2.3), for all $z\in\mathbf{D}(0,j_{\nu,1}')$ we obtain the inequality

$$\mathrm{Re}\left(1+\frac{zf_\nu''(z)}{f_\nu'(z)}\right)\geq1-\left(\frac{1}{\nu}-1\right)\sum_{n\geq1}\frac{2r^2}{j_{\nu,n}^2-r^2}-\sum_{n\geq1}\frac{2r^2}{j_{\nu,n}'^2-r^2},$$

where $|z|=r$. Moreover, observe that if we use the inequality (2.1) then we get that the above inequality is also valid when $\nu>1$. Here we used that the zeros $j_{\nu,n}$ and $j_{\nu,n}'$ interlace according to the inequalities

$$\nu\leq j_{\nu,1}'<j_{\nu,1}<j_{\nu,2}'<j_{\nu,2}<j_{\nu,3}'<\ldots.$$

Now, the above deduced inequality implies for $r \in (0, j'_{\nu,1})$

$$\inf_{z\in\mathbf{D}(0,r)}\left\{\operatorname{Re}\left(1+\frac{zf''_\nu(z)}{f'_\nu(z)}\right)\right\} = 1+\frac{rf''_\nu(r)}{f'_\nu(r)}.$$

On the other hand, the function $u_\nu : (0, j'_{\nu,1}) \to \mathbb{R}$, defined by

$$u_\nu(r) = 1+\frac{rf''_\nu(r)}{f'_\nu(r)},$$

is strictly decreasing since

$$\begin{aligned} u'_\nu(r) &= -\left(\frac{1}{\nu}-1\right)\sum_{n\geq1}\frac{4rj^2_{\nu,n}}{(j^2_{\nu,n}-r^2)^2} - \sum_{n\geq1}\frac{4rj'^2_{\nu,n}}{(j'^2_{\nu,n}-r^2)^2} \\ &< \sum_{n\geq1}\frac{4rj^2_{\nu,n}}{(j^2_{\nu,n}-r^2)^2} - \sum_{n\geq1}\frac{4rj'^2_{\nu,n}}{(j'^2_{\nu,n}-r^2)^2} < 0 \end{aligned}$$

for $\nu > 0$ and $r \in (0, j'_{\nu,1})$. Here we used again that the zeros $j_{\nu,n}$ and $j'_{\nu,n}$ interlace and for all $n \in \{1, 2, \dots\}$, $\nu > 0$ and $r < \sqrt{j_{\nu,1}j'_{\nu,1}}$ we have that

$$j^2_{\nu,n}(j'^2_{\nu,n}-r^2)^2 < j'^2_{\nu,n}(j^2_{\nu,n}-r^2)^2.$$

Observe also that $\lim_{r\searrow0} u_\nu(r) = 1 > \alpha$ and $\lim_{r\nearrow j'_{\nu,1}} u_\nu(r) = -\infty$, which means that for $z \in \mathbf{D}(0, r_1)$ we have

$$\operatorname{Re}\left(1+\frac{zf''_\nu(z)}{f'_\nu(z)}\right) > \alpha,$$

if and only if r_1 is the unique root of

$$1+\frac{rf''_\nu(r)}{f'_\nu(r)} = \alpha,$$

situated in $(0, j'_{\nu,1})$. □

Proof of Theorem 2. Observe that

$$z\frac{g''_\nu(z)}{g'_\nu(z)} = \frac{\nu(\nu-1)J_\nu(z)+2(1-\nu)zJ'_\nu(z)+z^2J''_\nu(z)}{(1-\nu)J_\nu(z)+zJ'_\nu(z)}.$$

The recurrence formulae $zJ'_\nu(z) = \nu J_\nu(z) - zJ_{\nu+1}(z)$ implies

$$z\frac{g''_\nu(z)}{g'_\nu(z)} = z\frac{zJ_{\nu+2}(z)-3J_{\nu+1}(z)}{J_\nu(z)-zJ_{\nu+1}(z)},$$

and using (2.4) it follows that

$$1+z\frac{g''_\nu(z)}{g'_\nu(z)} = 1-\sum_{n\geq1}\frac{2z^2}{\alpha^2_{\nu,n}-z^2}.$$

Application of the inequality (2.3) implies that

$$\operatorname{Re}\left(1+z\frac{g''_\nu(z)}{g'_\nu(z)}\right) \geq 1-\sum_{n\geq1}\frac{2r^2}{\alpha^2_{\nu,n}-r^2},$$

where $|z| = r$. Thus, for $r \in (0, \alpha_{\nu,1})$ we get

$$\inf_{z\in\mathbf{D}(0,r)}\left\{\operatorname{Re}\left(1+\frac{zg''_\nu(z)}{g'_\nu(z)}\right)\right\} = 1-\sum_{n\geq1}\frac{2r^2}{\alpha^2_{\nu,n}-r^2} = 1+\frac{rg''_\nu(r)}{g'_\nu(r)}.$$

The function $v_\nu : (0, \alpha_{\nu,1}) \to \mathbb{R}$, defined by

$$v_\nu(r) = 1 + \frac{r g''_\nu(r)}{g'_\nu(r)},$$

is strictly decreasing and

$$\lim_{r \searrow 0} v_\nu(r) = 1, \quad \lim_{r \nearrow \alpha_{\nu,1}} v_\nu(r) = -\infty.$$

Consequently, the equation

$$1 + \frac{r g''_\nu(r)}{g'_\nu(r)} = \alpha$$

has a unique root r_2 in $(0, \alpha_{\nu,1})$, and this equation is equivalent to (1.2). In other words, we have

$$\operatorname{Re}\left(1 + \frac{z g''_\nu(z)}{g'_\nu(z)}\right) > \alpha,\ z \in \mathbf{D}(0, r_2) \ \text{ and } \inf_{z \in \mathbf{D}(0,r_2)} \operatorname{Re}\left(1 + \frac{z g''_\nu(z)}{g'_\nu(z)}\right) = \alpha.$$

Finally, let us recall that (see [**Wa**, p. 597]) when $\rho+\nu > 0$ and $\nu > -1$ the so-called Dini function $z \mapsto zJ'_\nu(z) + \rho J_\nu(z)$ has only real zeros and according to Ismail and Muldoon [**IM**, p. 11] we know that the smallest positive zero of the above function is less than $j_{\nu,1}$. This in turn implies that $\alpha_{\nu,1} < j_{\nu,1}$, which completes the proof. $\square$

References

[**Ba1**] Á. Baricz, Geometric properties of generalized Bessel functions of complex order, *Mathematica* 48(71) (2006) 13–18.

[**Ba2**] Á. Baricz, Geometric properties of generalized Bessel functions, *Publ. Math. Debrecen* 73 (2008) 155–178.

[**Ba3**] Á. Baricz, *Generalized Bessel Functions of the First Kind*, Lecture Notes in Mathematics, vol. 1994, Springer-Verlag, Berlin, 2010.

[**BKS**] Á. Baricz, P.A. Kupán, R. Szász, The radius of starlikeness of normalized Bessel functions of the first kind, *Proc. Amer. Math. Soc.* (in press).

[**BP**] Á. Baricz, S. Ponnusamy, Starlikeness and convexity of generalized Bessel functions, *Integr. Transforms Spec. Funct.* 21 (2010) 641-653.

[**Br**] R.K. Brown, Univalence of Bessel functions, *Proc. Amer. Math. Soc.* 11(2) (1960) 278–283.

[**IM**] M.E.H. Ismail, M.E. Muldoon, Bounds for the small real and purely imaginary zeros of Bessel and related functions, *Methods Appl. Anal.* 2(1) (1995) 1–21.

[**KT**] E. Kreyszig, J. Todd, The radius of univalence of Bessel functions, *Illinois J. Math.* 4 (1960) 143–149.

[**OLBC**] F.W.J. Olver, D.W. Lozier, R.F. Boisvert, C.W. Clark (Eds.), *NIST Handbook of Mathematical Functions*, Cambridge Univ. Press, Cambridge, 2010.

[**Sz**] R. Szász, On starlikeness of Bessel functions of the first kind, In: Proceedings of the 8th Joint Conference on Mathematics and Computer Science, Komárno, Slovakia, 2010, 9pp.

[**SK**] R. Szász, P.A. Kupán, About the univalence of the Bessel functions, *Stud. Univ. Babeş-Bolyai Math.* 54(1) (2009) 127–132.

[**Wa**] G.N. Watson, *A Treatise of the Theory of Bessel Functions*, Cambridge University Press, Cambridge, 1995.

Department of Economics, Babeş-Bolyai University, 400591 Cluj-Napoca, Romania
E-mail address: `bariczocsi@yahoo.com`

Department of Mathematics and Informatics, Sapientia Hungarian University of Transylvania, 540485 Târgu-Mureş, Romania
E-mail address: `rszasz@ms.sapientia.ro`